Research Article

Agronomic Performance and Genotype $\times$ Environment Interaction of some Hybrid Rice in Nigeria

Bashir, M., Ehirim, B.O, Isong, A., Ejoro, O., Odoba, A., Hadiza, A.B. and Aliyu, U.

National Cereals Research Institute, Badeggi. P.M.B, 8, Bida. Nigeria

Corresponding Author E-mail: mbashiru00@gmail.com

Abstract

Agronomic parameters and Genotype $\times$ environmental interaction for stability estimate were investigated on grain yield of 2 hybrid rice lines (TEJ Gold and 6444 Gold) with three commercially released rice varieties (FARO 44, FARO 52 and FARO 57) as checks. This was done in six locations in 2017 of which five out of the six (Kebbi, Kano, Enugu, Warri and Lafiagi) were in dry season and one (Wushishi) in wet season. The experiments were laid out in a randomized complete block design in three replications having 20 cm inter and intra row spacing. AMMI ANOVA for grain yield revealed significant difference among the genotypes and environments that were tested ($P < 0.01$) and their interaction. The observed difference on the interaction indicated that, the genotypes responded differently across the different environments. The partitioning of GGE through GGE biplot analysis showed that, PC1 and PC2 contributed 75.34% and 16.33% which summed up and explained a total of 91.67% of the observed variation seen, though PC1 contributed more of the observable variations. Cultivar ranking and genotype stability revealed that the hybrid TEJ GOLD, performed well across the test locations though not stable across the locations tested. However, FARO 52 and FARO 57 had a small GE effect, indicating their stability and being less influenced by the environment. The two hybrids rice (TEJ Gold and 6444 Gold) significantly gave a higher yield of 6683 and 6005 kg/ha respectively, with yield advantage ranging from 11.9 to 32.8% over the three check varieties.

Keywords: Agronomic performance, Genotype, Grain yield, Hybrid, Interaction, Nigeria

© 2019 National Cereals Research Institute (NCRI), Nigeria, all rights reserved.

Introduction

Rice has become an important strategic and daily staple food crop in Nigeria. The potential land area for rice production in Nigeria is between 4.6 million and 4.9 million hectares (FAO, 2006). Out of this, only about 1.7 million ha or 35 percent of the available land area is presently under rice cultivation (Olaf *et al.*, 2003). There is need to produce more rice in Nigeria to meet demands created by increasing population, mass exodus of rural people to urban area, effect of climate change and rising incomes cannot be overemphasized. This production increase must be achieved on less land, with less labour, less water, and less pesticides, and must be sustainable. (Rajesh *et al.*, 2012) Small-scale farmers with farm holdings of less than 1 hectare cultivate most of the rice produced in Nigeria (Imolehin *et al.*, 2000). However, rice productivity and production

at the farm level are constrained by several factors. These factors include insufficient and appropriate technologies, biotic factors, poor supply of inputs, ineffective farmer organizations and groups, low yield and poor milling quality of local rice varieties, poor marketing arrangements, inconsistent agricultural input and rice trade policies, poor extension systems and environmental constraints (Kueneman, 2006). These environmental constraints include poor drainage and iron toxicity in undeveloped lowland swamps, drought, deficiencies in N and P, insufficient rainfall and poor management practices (Janaiah, 2000). Experience in China, India, and Vietnam have established that hybrid rice offers an economically viable option to increasing varietal yields beyond the present level of 1 to 2.5 tons/ha of rice paddy. Several other countries such as Bangladesh, Brazil, Colombia,

Egypt, Democratic People's Republic of Korea, Japan, Malaysia, Myanmar, Pakistan, Philippines, Republic of Korea, Sri Lanka, Thailand and USA are currently exploring the prospects of hybrid rice. Availability of adequately trained human resources is an essential prerequisite for developing an effective national hybrid rice breeding program. (Rajesh *et al.*, 2012)

Hybrid rice breeding uses several concepts, skills, and procedures which are strikingly different from those used for inbred rice breeding. These must be learned by plant breeders before initiating a comprehensive hybrid rice breeding program (Yuan, 2002). Hybrid rice is a proven and successful technology for rice production and can contribute significantly toward improving food security, raising rice productivity and farmers' income, and providing employment opportunities (George, 2007). Remarkable progress continues as hybrid rice technology makes its way across Asia and to other countries (IRRI, 1994, IRRI, 1996). Since the 4th International Hybrid Rice Symposium held in Hanoi, Vietnam, in 2002, the area grown to hybrid rice worldwide increased to an estimated 19.8 million hectares in 2007, including about 2.8 million hectares outside China but no area cultivated in Nigeria till date. All major rice-producing countries in the world have been investing in applying hybrid rice technology, and, in recent years, the seed industry has also been involved in hybrid rice research and development (Mao, 2001). In this research paper, two hybrid rice genotypes were evaluated in six locations to assess their stability and yield potential.

Materials and Methods

Two hybrid rice genotypes (6444 GOLD and TEJ GOLD), with three inbred checks (FAO44 FARO 52 and FARO 57) were evaluated in 2017 cropping season (wet and dry season) in six locations. Kebbi, Kano, Enugu, Warri and Lafiagi were evaluated during dry season of 2017 while Wushishi location was evaluated in the wet season of same year, all in a Randomized Complete Block design with three replications respectively. Each entry was planted in a 4m x 3m plot size and fertilizer application was at 80, 40, 40 kg per ha of N, P₂O₅ and K₂O, with N applied at two splits at 21 and 42 days after transplanting (DAT).

Weeding was controlled by propanil and 2-4-D formulation at 4 liters per ha of Orizo Plus, with supplementary hand weeding. Agronomic data were collected for plant height (cm), Number of tillers per square meter, Number of Panicles per square meter, Number of days to 50% flowering and grain yield. This was subjected to analysis with CropStat, version 3.0. Stability and Sensitivity on the rice grain yield was determined by subjecting the grain yield data to additive main effect and multiplicative interaction (AMMI), GGE-biplot and Boxplot analysis using Breeding View (BV) statistical package in the breeding management system, version 3.0.9.

Results and Discussions:

Table 1 shows that plant height in all the locations investigated were significantly different. FARO 57 had the highest mean plant height of 113.1 cm while FARO 44 had the lowest mean plant height of 92.1cm. However, hybrid rice TEJ GOLD had a mean plant height of 98.9 cm which is within the range of plant height accepted by local farmers in Nigeria. Looking at the specific locations for plant height, FARO 57 in Kebbi location had the highest plant height value of 138.4 and FARO 44 in Enugu location and had the lowest plant height value. It is also important to note that genotypes with plant height figures of about 120 cm could lead to lodging. Tiller number /m² in table 1 showed that Kano, Lafiagi and Kebbi locations differed significantly for number of tillers per square meter while Warri Enugu and Wushishi were not significant. Wushishi location recorded more tillers when compared to other locations where the entries were tested with hybrid 6444GOLD recording the highest number of tillers (738 tillers /m²). However, FARO 44 had the highest mean tiller number (535 tillers/m²) when compared across the locations investigated. Number of panicles/m² in Table 2 showed that Kano, Lafiagi and Kebbi showed wide range of difference for number of panicles/m². Warri, Enugu and Wushishi were not significantly different for number of panicles/m². For the 6 locations that were used for the investigation, Wushishi location showed to have the highest number of panicles with 6444 GOLD recording the highest (432) though there was no significant difference observed among the 5 test entries. On the other hand, number of days to 50% flowering

showed significant difference for the 6 location investigated. Though the mean days to 50% flowering showed that hybrid 6444 GOLD recorded the highest (93 days), Lafiagi showed to have the latest number of days to 50% flowering while Wushishi showed to be the earliest.

Grain yield data in Table 3 shows that the test entries investigated differed significantly in Kano, Lafiagi, Kebbi, Enugu and Wushishi but was not significantly different at Warri location. Warri location recorded the least grain yield when compared to other locations. However, TEJ GOLD recorded the highest mean grain yield of 6.683 metric tons per hectare. 6444 GOLD also recorded the highest grain yield of 9.967 tons per hectare in wushishi location. It was also observed that the two hybrid test entries outperformed the check varieties in all the 6 locations. 6444 GOLD had a 19.3, 10.6 and 11.9% yield advantage over FARO 57, FARO 52 and FARO 44 respectively. TEJ GOLD had a 32.2%, 23.1% and 24.6% yield advantage over FARO 57, FARO 52 and FARO 44 respectively. It could also be inferred that the high yield experienced in Wushishi location may be as a result of the experiment being done in the wet season of 2017. From the Finlay and Wilkinson regression analysis and ANOVA (Table 4), significant differences were observed among the test genotype across the environment which shows the wide differences in the varieties used for the experiment as well as their response to the test environments. The test environments on the other hand differed significantly also showing that the quality of the test environment were not the same from location to location as shown in Table 4. IPCA 1 and IPCA 2 were significantly different explaining the observable difference in the behavior of the genotypes in their respective environments. Identifying high yielding and stable genotypes is easier when such significant differences are observed (Eberhart and Russell, 1966; Ishaq *et al.*, 2015).

Table 5 shows the mean values for yield and regression coefficient (b), for 2 hybrid rice varieties over four environments. Genotypic sensitivity which shows the changes in the environmental quality is shown by the slope (b value). This explains the changes experienced in

the quality of environments; where values of $b > 1$ mean genotypes with a higher sensitivity and less stability than average sensitivity ($b=1$), while $b < 1$ means genotypes that were less sensitive and more stable. Therefore, the best genotypes within the seven (7) environments of study were those within the $b = <1$ range of sensitivity and should be noted that the lesser that b Value the more stable the variety. From Table 5, the sensitivity ranking shows that FARO 52 was the most stable variety across the locations (0.6096) though its mean yield indicated that it ranked 4th place in cultivar superiority with a mean grain yield of 5431tons/ha. However, hybrid cultivar TEJ GOLD, had the highest mean grain yield of 6683tons/ha as shown from the cultivar ranking score of 1 but it showed to be very sensitive as it scored 4th in the sensitivity ranking (1.1688). It could be deduced that, though the hybrid test entry TEJ GOLD variety outperformed the 3 check varieties in terms of yield, it is however, not a better variety in terms of stability across the ranges of environment. This finding agrees with Bashir *et al.* (2018) and Maji *et al.* (2015) who opined that, an ideal genotype is the genotype with high performance combined with good stability to different environments. Figure 1 shows AMMI (additive means and multiplicative interactions) biplots and environmental scaling of the 5 test entries in 6 environments. Three mega environments were described from Figure 1. Lafiagi and Kano formed different mega environment while the others formed one cluster showing to have almost same environmental quality. The partitioning of GGE through GGE biplot analysis of grain yield showed that, PC1 and PC2 explained 75.34% and 16.33% of GGE sum of squares respectively, explaining a total of 91.67% of the observed variation seen though, PC1 contributed more. However, TEJ GOLD being on the vertex hexagon point of the upper right side of the GGE environmental scaling makes it the most stable and best genotype across the locations investigated.

Conclusion

Agronomic performance of crop varieties is a sure way of distinguishing varietal differences and a good guide for breeders when making selection. AMMI statistical model is a tool in selecting the most suitable and stable high yielding crop

genotype for specific as well as for diverse environments. In the present study, both agronomic performance and AMMI model has shown that, Hybrid TEJ GOLD performed well across the test locations though, shown not stable across the locations tested. The largest proportion of the total variation observed in the grain yield of the rice genotypes is attributed to environments. The mean grain yield value of genotypes averaged over environments indicated that FARO 52 and FARO 57 had a small GE effect, indicating their stability and being less influenced by the environment. The 2 hybrid test entries should be re-evaluated across many environments and tested for stability before recommendation for release.

References

- Bashir, M., Ehirim, B. O., Maji, A.T., Bakare, S. O., Isong, A. and Odoba, A. (2018). Evaluation of Yield Components and Genotype $\times$ Environment Interaction on Grain Yield of Submergence Lowland Rice Genotypes, *Int. J. Pure App. Biosci.* 6(3), 623-630
- Breeding Management System (BMS). Version 3.0.9, *Integrated Breeding Platform* (IBP), Mexico (2015)
- Eberhart, S.A and Russell, W. L. (1966). Stability parameters for comparing varieties. *Crop Sci.* 6, 36-40.
- FAO (2006). World agriculture: towards 2030/2050. Interim report. Rome (Italy): FAO.
- Finlay, K., Wilkinson, G. (1963). The analysis of adaptation in a plant-breeding programme. *Crop Pasture Sci.* 14, 742-754
- George, A. (2007). Principles of Plant Genetics and Breeding; BLACKWELL PUBLISHING: ISBN-13: 978-1-4051-3646-4 pp 15-583
- Ishaq, M. N., Agrama, H. and Adeleke, A., (2015). Exploiting genotype $\times$ environment interaction in soybean breeding in Nigeria. *Int. J. Adv. Res. Biol. Sci.* 2(1), 24-32
- Imolehin, E.D. and Wada, A.C. (2000). Meeting the rice production and consumption demands of Nigeria with improved technologies. *International Rice Commission Newsletter.* 49, FAO, Rome, Pp. 23-41.
- IRRI (1996). "Hybrid Rice". Proc. of the International Symposium on Hybrid rice, 6-10 October, 1996, Changsha, Hunan, China, International Rice Research Institute, Manila Philippines. Pp. 201-313
- IRRI (1994). "Hybrid Rice Technology- New Development and Future prospects." Ed. S. S. Virmani. International Rice Research Institute, Manila.
- Janaiah, A. and Hossain, M. (2000). Hybrid rice for food security in the tropics: An evaluation of farm-level experiences in India. Paper presented at the 3rd International Crop Science Congress held in Hamburg, Germany, 17-22 August
- Kueneman E.A. (2006). Improved rice production in a changing environment: from concept to practice. *IRC News* 1. 55, 1-20.
- Mao, C.X. (2001). Improving seed production to speed up the global commercialization by hybrid rice. *Rice Research for Food Security and Poverty Alleviation*, S. Peng and B. Hardy (Eds.). Proceedings of the International Rice Research Conference, 31 March-3 April 2000. International Rice Research Institute, Los Banos, Laguna, Philippines. pp. 692
- Maji, A. T., Bashir, M., Odoba, A., Gbanguba, A. U. and Audu, S. D. (2015). Genotype $\times$ Environment Interaction and Stability Estimate for Grain Yield of Upland Rice Genotypes in Nigeria. *J Rice Res.* 3, 136.
- Olaf, E., Frederic, L., Akande, S.O., Titilola, Akpokodje G. and Ogundele, O. O. (2003). The Nigerian Rice Economy in A Competitive World: Constraints, Opportunities and Strategic Choices. *Rice production systems in Nigeria: A survey* 8-73
- Rajesh, S. and Lekha, R. (2012) Ideal Hybrid Rice Seed Production Package: An Overview. *Indian Research Journal of Extension Education, Special Issue, 2*, 244-251
- Yuan L.P. (2002). Recent progress in the development of hybrid rice in China. In: Adoption of hybrid rice in Asia: policy support. Rome, FAO. 65-68.

Table 1. Plant height and Number of Tillers per square meter in Multi-Locational evaluation of Hybrid rice lines and check varieties

Designation	Plant Height (cm)							Number of Tiller/m ²						
	Kano	Lafiagi	Kebbi	Warri	Enugu	Wushishi	Mean	Kano	Lafiagi	Kebbi	Warri	Enugu	Wushishi	Mean
6444 GOLD	113.0	104.3	114.5	90.9	95.9	120.0	104.5	590	350	633	543	270	738	531
TEJ GOLD	97.3	104.7	108.3	78.5	91.7	114.0	98.9	530	445	617	458	298	683	524
FARO 44 ©	91.7	94.6	99.4	73.5	82.8	110.3	92.1	608	432	508	533	337	682	535
FARO 52 ©	101.3	126.6	121.5	97.6	102.9	120.0	110.8	517	422	550	585	250	537	483
FARO 57 ©	106.7	111.3	138.4	91.7	101.0	132.3	113.1	520	323	542	542	250	590	468
Lsd 5%	*	*	*	*	*	*	*	*	*	*	Ns	Ns	Ns	*
CV %	2.5	3.3	5.0	4.3	4.3	4.9	3.3	2.9	4.3	6.6	12.8	15.3	17.7	12.0

*-Significant, ns – Not Significant, © - check entry

Table 2. Number of Panicle per square meter and Number of days to 50% flowering in Multi-Locational evaluation of Hybrid rice lines and check varieties

Designation	Number of Panicles/m ²							Number of Days to 50% Flowering					
	Kano	Lafiagi	Kebbi	Warri	Enugu	Wushishi	Mean	Kano	Lafiagi	Warri	Enugu	Wushishi	Mean
6444 GOLD	300	333	562	377	492	620	432	74	109	89	108	68	93
TEJ GOLD	296	297	525	332	540	610	415	71	85	80	92	58	80
FARO 44©	307	353	418	412	552	557	421	70	100	86	98	61	86
FARO 52©	225	253	503	350	444	432	358	72	105	103	99	69	92
FARO 57©	225	313	437	350	432	453	360	74	108	89	102	67	91
Lsd 5%	*	*	*	ns	ns	ns	*	*	*	*	*	*	*
CV %	1.9	5.0	9.4	21.2	12.1	17.0	12.3	0.5	2.5	5.2	2.5	1.1	1.5

*-Significant, ns – Not Significant, © - check entry

Table 3. Grain yield and Percentage yield advantage over checks Hybrid rice lines in Multi-Locational evaluation

Designation	Grain Yield (kg/ha)							% Yield Adv over checks		
	Kano	Lafiagi	Kebbi	Warri	Enugu	Wushishi	Mean	FARO 57	FARO 52	FARO44
6444 GOLD	7267	4467	6933	4100	4200	9967	6005	19.3	10.6	11.9
TEJ GOLD	7300	5733	7733	3917	6067	9767	6683	32.8	23.1	24.6
FARO 44©	7333	4167	6567	3550	4467	7267	5364			
FARO 52©	6000	6067	6533	3450	3667	6167	5431			
FARO 57©	6900	4100	6333	3367	4167	5733	5033			
Lsd 5%	*	*	*	ns	*	*	*			
CV %	6.6	8.0	4.9	16.2	7.0	11.6	9.7			

*-Significant, ns – Not Significant, © - check entry

Table 4. Analysis of variance using F and W Regression Analysis

Source	D.F.	S.S.	M.S.	V.R.	F Pr
Genotypes	4	11826423	2956606	4.19	0.0104
Environments	5	67432941	11238823	15.91	<0.001
Interactions	24	16953257	706386		
IPCA 1	9	10968241	1218693	6.35	0.0080
IPCA 2	7	4450130	635733	3.31	0.0575
Residuals	8	1534886	191861		

Table 5. Ranking and Genotype stability Parameters for grain yield based on the Finlay and Wilkinson (1963) stability model.

Genotype	Mean	S.E.	Sensitivity (b)	Sensitivity Ranking	Cultivar Superiority	Cultivar Ranking	Mean square Deviation
6444 GOLD	6005	293.2	1.3957	5	13295	2	360557
FARO 44	5364	293.2	1.0344	3	575138	3	263479
FARO 52	5431	293.2	0.6096	1	1385325	4	994633
FARO 57	5033	293.2	0.7609	2	1704096	5	473710
TEJ GOLD	6683	293.2	1.1688	4	2197184	1	314276

Figure 1. GGE biplot for best rice genotypes in different environment for grain yield