Available online: www.ncribjare.org
 ISSN: 2695-2122, e-ISSN: 2695-2114
 DOI: <https://doi.org/10.35849/BJARE202102011>
 Journal homepage: www.ncribjare.org

Research Article

Effects of Ginger and Garlic on the Proximate and Mineral Compositions of Differently Processed Nile Tilapia (*Oreochromis niloticus*)

Gana, A. B.¹, Yusuf, D. U.¹, Ibrahim, R.¹, Bake, G. G.², Iriobe, T.¹, Jega, I. S.¹ and Abubakar, F. N.²

¹Department of Forestry and Fisheries, Faculty of Agriculture, Kebbi State University of Science and Technology, Aliero, Kebbi State, Nigeria.

²Department of Water Resources, Aquaculture and Fisheries Technology, School of Agriculture and Agricultural Technology, Federal University of Technology, Minna, Niger State, Nigeria.

Corresponding e-mail: abglatiko@gmail.com

Abstract

Freshly caught Nile Tilapia (*Oreochromis niloticus*) samples were obtained from Sabiyel Lake in Aliero Local Government Area of Kebbi State. *Oreochromis niloticus* samples were descaled, degutted, cut into chunks and rinsed with clean water at Fisheries Laboratory, Department of Forestry and Fisheries, Kebbi State University of Science and Technology, Aliero. The samples were subjected to different treatments designated as; T1 (oven drying), T2 (Frying), T3 (smoking), T4 (oven-dried + 5% ginger-garlic), T5 (Fried + 5% ginger-garlic), T6 (smoked + 5% ginger-garlic), T7 (oven-dried + 10% ginger-garlic), T8 (Fried + 10% ginger-garlic), T9 (smoked + 10% ginger-garlic), T10 (oven-dried + 15% ginger-garlic), T11 (Fried + 15% ginger-garlic), T12 (smoked + 15% ginger-garlic). The processed samples were taken to Agric. Chemical Laboratory, Faculty of Agriculture, Usmanu Danfodio University Sokoto, Nigeria, for the determination of proximate and mineral compositions. The treatments were analysed in 2 phases: at week 0 for the first phase of analysis and after 8 weeks of storage in an airtight plastic container at room temperature. The samples were then taken to the laboratory for the second phase of analysis. This study revealed that processing (oven drying, frying and smoking) of *Oreochromis niloticus* with ginger-garlic gave high nutritive content after 8 weeks of storage and thus might prevent the use of obnoxious substances as a means of preservation and substantially improve consumer's satisfaction and quality fish protein intake.

Keywords: Ginger, Garlic, *Oreochromis niloticus*, Proximate, Mineral contents

© 2021 National Cereals Research Institute (NCRI), Nigeria, all rights reserved.

Introduction

Fish is an excellent source of good quality protein containing all the essential amino acids in desirable concentrations for human consumption (Omitoyin, 2007). It is eaten fresh or processed and forms a much-cherished delicacy that cut across socioeconomic, ages, religious and educational barriers (Adebayo-Tayo *et al.*, 2008). In Nigeria, a greater portion of fish caught are preserved by various processing methods and only a negligible proportion are marketed fresh (Onyia and Adebayo, 2008). Fish processing is aimed at preventing or reducing post-harvest losses due to

mould, bacteria, and insect's infestation (Abolagba and Nuntah, 2011). The processed products have proved relevant to inhibiting microbial growth, improving acceptability, enhancing flavour and above all prolonging shelf life of the products either by way of refrigerating, salt curing, drying, frying, boiling, smoking or use of preservatives (Eyo, 2001; Yakubu and Ngueku, 2015). Unfortunately, many fish sellers use poisonous substances (dichlorvos, lindane, snipper or *pia pia*) to halt insects or microbial infestation and not minding the health implications of the consumers (Musa *et al.*, 2010). However, Idris *et al.* (2010)

reported that the dipping of fish in a concentration of preservatives (ginger) before smoking has beneficial effects on the overall quality of the products.

Ginger is a complex substance comprising many compounds (Srivastava *et al.*, 2000). The essential oil and resin collectively known as oleoresin are contained in the ginger rhizome. However, this essential oil varies according to geographical origins. Akoachere *et al.* (2002); Young *et al.* (2005) and Al-Awwadi (2017) revealed that ginger is a potential herb used worldwide for its immense phytotherapeutic properties (antimicrobial, antibiotic and antioxidant effects, an ability to inhibit the formation of inflammatory compounds and direct anti-inflammatory effects). It is also effective against some kind of cancers (Park *et al.*, 2006), controls blood pressure and hypertension (Ajay *et al.*, 2003; Ranilla *et al.*, 2010), stimulates blood circulation, helps in lowering cholesterol and is associated with combating heart problems. Ginger and some other herbs are used as a food preservative and can increase food safety and shelf life of fatty and processed food products (El-Baroty *et al.*, 2010).

Garlic is considered to be one of the best disease preventive foods because of its potent and widespread effects (Amagase *et al.*, 2001; Papu *et al.*, 2017). Many favourable experimental and clinical effects of the consumption of garlic preparations, including garlic oil, garlic extracts, and garlic powder with differences in chemical compositions, have been reported (Lanzotti *et al.*, 2014). These effects have been substantiated by modern research and garlic has been proved to elicit antimicrobial (Yoshida *et al.*, 1998), antihypertensive (Suetsuna, 1998), hepatoprotective (Wang *et al.*, 1998), hypolipidemic (Sumiyoshi, 1997), antidiabetic (Ahmed and Sharma, 1997; Ohaeri and Adoga, 1998) and insecticidal (Singh *et al.*, 1996) properties. The biological responses include reduction of risk factors for cardiovascular diseases and cancer, stimulation of immune function, enhanced foreign compound detoxification, radio-protection, restoration of physical strength, resistance to various stresses and potential antiaging effects (Amagase *et al.*, 2001). Garlic is famous for its characteristic odour,

arising from allicin and other oil-soluble sulphur components (Amagase, 2006). Processed garlic contains a variety of sulphur-containing compounds other than those found naturally in intact garlic cloves, depending on the conditions applied (Fenwick and Hanley, 1985; Lawson, 1993; Pingxiang *et al.*, 2020). However, the sulphur-containing compounds in garlic preparation considerably vary, depending on their methods of processing (Amagase *et al.*, 2001). Although garlic has been used safely in cooking as a popular condiment or flavouring and used traditionally for medicinal purposes, also it is commonly known that excessive consumption of garlic can cause a problem (Mader, 1990; Siegers, 1992).

Ginger and garlic can inhibit and kill bacteria, fungi, lower blood pressure, cholesterol and sugar, also prevent blood clotting, and contains anti-tumour properties (Gebreselema and Mebrahtu, 2013; Chinedu and Jivini, 2019). It can also boost the immune system to fight off potential disease and maintain health (Abdullah *et al.*, 1988; Rodrigo *et al.*, 2015). Therefore, this study is aimed at investigating the effects of ginger and garlic on the proximate and mineral compositions of differently processed *Oreochromis niloticus*.

Materials and Methods

Description of the study area

The experiment was conducted in the Fisheries Laboratory, Department of Forestry and Fisheries, Kebbi State University of Science and Technology Aliero, Kebbi State Nigeria. Aliero is located in the Southeast of Kebbi State between approximate longitudes 3°6'E to 4°27'E and latitudes 11°03'N to 12°47'N (Singh, 2013; Ibrahim *et al.*, 2018).

Ginger and garlic preparation

The ginger and garlic were purchased from Aliero market, in Aliero Local Government Area of Kebbi State. The samples were rinsed with clean water, sun-dried, separately ground into a powder using mortar and pestle, then packed separately in an airtight plastic container, and stored till when needed.

Fish sample identification and collection

The fish sample was identified using a field guide by Olaosebikan and Raji (2004). Freshly harvested

Nile Tilapia (*Oreochromis niloticus*) were collected at the landing site of Sabiyel Lake in Aliero Local Government Area of Kebbi State, Nigeria and transported to the Fisheries Laboratory in a plastic flask containing ice blocks.

Preparation of fish sample

Oreochromis niloticus samples were descaled, degutted, cut into chunks and rinse with clean water at Fisheries Laboratory, Department of Forestry and Fisheries, Kebbi State University of Science and Technology, Aliero. Zero (0) grams (0%: Control), 50 grams (5%), 100 grams (10%) and 150 grams (15%) of ginger and garlic (1:1) per kilogram of *Oreochromis niloticus* were measured using a sensitive weighing balance (Tanita KD-160: model). The weighed ginger and garlic were mixed in a clean plastic bowl, containing 100 ml of clean water, the sample fish were then placed inside for 20 minutes. The fish sample was then subjected to different processing methods (oven-drying, frying and smoking).

Chemical analysis

The processed samples were taken to Agric. Chemical Laboratory, Faculty of Agriculture, Usmanu Danfodio University Sokoto, Nigeria, for the determination of chemical compositions [moisture, ether extract, crude fibre, ash content, crude protein and nitrogen-free extract (NFE)], and mineral compositions (sodium, phosphorus, potassium, magnesium, and calcium). The proximate and mineral compositions of the treatments were determined using the procedure described in Gana *et al.* (2020). The treatments were analysed in 2 phases, at week 0 for the first phase of analysis and after 8 weeks of storage in an airtight plastic container at room temperature, the samples were then taken to the laboratory for the second phase of analysis.

Experimental design

The samples were subjected to different treatment and designated as; T1 (oven drying), T2 (Frying), T3 (smoking), T4 (oven dried + 5% ginger-garlic), T5 (Fried + 5% ginger-garlic), T6 (smoked + 5% ginger-garlic), T7 (oven dried + 10% ginger-garlic), T8 (Fried + 10% ginger-garlic), T9 (smoked + 10% ginger-garlic), T10 (oven dried + 15% ginger-garlic), T11 (Fried + 15% ginger-

garlic), T12 (smoked + 15% ginger-garlic). Each treatment was replicated three times.

Data analysis

The collected data were subjected to analysis of variance (ANOVA) and where there was a significant difference, Duncan's Multiple Range Test (DMRT) was applied to rank the treatment means ($P < 0.05$). All statistical analyses were computed using MINITAB (V14) statistical package for Windows.

Results and Discussion

Proximate and mineral compositions of ginger and garlic

The proximate analysis of ginger and garlic revealed (Table 1) ginger had 5.6%, 4.50% and 7.50% while garlic had 9.19% 6.00% and 6.50% crude protein, moisture content and ether extract, respectively. Garlic contains 3.07 mg/kg, 1.45 mg/kg and 85.00 mg/kg while ginger contains 3.64 mg/kg, 1.60 mg/kg and 70.00 mg/kg of phosphorus, magnesium and sodium, respectively. This result differs from the findings of Nwinuka *et al.* (2005) and Otunola *et al.* (2010). The variations may be attributed to processing methods and geographical locations.

Proximate compositions of differently processed *Oreochromis niloticus* at week zero and after eight weeks of storage

Significant changes in the nutrient values of *Oreochromis niloticus* subjected to different processing methods (oven drying, frying and smoking) at different ginger-garlic percentages were observed. Stephen *et al.* (2010); Adeyemi *et al.* (2013); Aberoumand (2014) reported that, processing compromises the nutritional quality of fish. The result (Table 2) showed that there is no significant difference ($P \geq 0.05$) between T2, T8 and T9 in ether extracts. However, there were significant differences ($P \leq 0.05$) among the treatment means in ether extract after eight weeks of storage. The higher ether extract was recorded by frying at week zero (0) and also at week (8) compared to others (oven drying and smoking). This could be attributed to oil absorption during the frying process (El-Lahamy *et al.*, 2019). Frying improves the sensory quality of food by the formation of aroma compounds texture and attractive colour (Aberoumand and Ziaei-Nejad,

2015). The application of suitable substances that has both antioxidant and antimicrobial activities may be useful to extend the shelf life and prevent economic loss (Yin and Chang, 2003). Ginger and garlic possess anti-oxidant, antiseptic and bacteriostatic properties (Iheagwara, 2013) and they are added to food to delay the onset of deterioration such as rancidity (Abdel-Hamied *et al.*, 2009). The T8 had the highest crude protein at week 0, though it was not significantly different from T2, T9 and T11. This could be attributed to different percentages of ginger-garlic to the treatments, resulting from their anti-microbial property. However, after 8 weeks of storage, the T1 (oven drying) had the highest crude protein and differ significantly among the treatment means. Chukwu and Shaba (2009) reported that oven drying could improve protein quality and prevent lipid oxidation in catfish. The higher protein content could be due to subsequent dehydration of the fish samples during storage (Synnes *et al.*, 2007). The T1 (oven drying) had the higher moisture contents at week 0 while T6 (smoked + 5% ginger-garlic) had the highest moisture content after 8 weeks of storage. The higher moisture content may be attributed to a rise in microbial activities during storage (Hassan and Umar, 2004). The moisture content decreases in all the fried samples. This could be attributed to moisture loss at the time of frying (Zafar *et al.*, 2019). Dashak *et al.* (2001) reported that reduced moisture content is an indication that the shelf life would be prolonged. Sampaio *et al.* (2009) reported that the determination of moisture content is one of the most common tests in foods since the water content in foods has an important relationship between conservation and the chemical, physical and microbiological changes during the storage.

Mineral composition of differently processed *Oreochromis niloticus* at week zero and after eight weeks of storage

The result (Table 3) revealed that T7 and T12 at week 0 were significantly different ($P \leq 0.05$) among the treatments in sodium. However, T12 (smoked + 15% ginger-garlic) was higher. Olopade *et al.* (2013) reported that duration, air humidity and temperature affect the efficiency of smoking and the quality of the final product. While after 8 weeks of storage, T1 had the highest sodium content and significantly differ among the

treatment means. At week 0, T4, T7 and T10 had the lower calcium content while after week 8, T1 had the highest value and was significantly different from other treatment means. Also, at week 0, all the treatments values recorded were higher compared to values recorded after 8 weeks of storage. This may be as a result of storage and possibly the addition of ginger-garlic. However, low mineral contents were observed in the treatments subjected to frying. This might be attributed to the fact that frying extracted some minerals into the fat (Hossein *et al.*, 2014; Karimian-Khosroshahi *et al.*, 2016).

Conclusion

This study concludes that *Oreochromis niloticus* has a high nutritive value. The methods of processing (oven drying, frying and smoking) have a significant effect on the proximate and mineral contents of the samples. Therefore, the use of ginger-garlic up to 15 % in the processing of *Oreochromis niloticus* might prevent the use of obnoxious substances as a means of preservation and substantially improve consumer's satisfaction and quality fish protein intake.

References

- Abdel-Hamied, A. A., Nassar, A. G. and El-Badry, N. (2009). Investigations of antioxidant and antibacterial activities of some natural extracts. *World Journal of Dairy and Food Sciences*, 4: 1 – 7.
- Abdullah, T. H., Kandil, O., Elkadi, A. and Carter, J. (1988). Garlic revisited: Therapeutic for the major diseases of our times. *Journal of the National Medical Association*, 80: 439 – 445.
- Aberoumand, A. (2014). Nutrient composition analysis of gish fish fillets affected by different cooking methods. *International Food Research Journal*, 21(5): 1989 – 1991.
- Aberoumand, A. and Ziaei-Nejad, S. (2015). Effect of cooking on quality commonly consumed marine fish platycephalidae (*Platycephalus indicus*) in Iran. *Turkish Journal of Agriculture - Food Science and Technology*, 3(11): 891 – 893.
- Abolagba, O. J., and Nuntah, J. N. (2011). Processing and distribution of smoked *Clarias* spp. in Benin City, Edo

- State. *International Research Journal of Biotechnology*, **2**(9), 213-219.
- Adebayo – Tayo, D. A., Agolabi, O. and Omosola, A. (2008). Quality changes of Nigerian Traditionally Processed fish species. Nature and Organoleptic changes *Journal of Food Technology*, **19**: 333-334.
- Adeyemi, O. T., Osilesi, O. O., Onajobi, F., Adebawo, O., Oyedemi, S. O. and Afolayan, A. J. (2013). Effect of processing on the proximate and mineral compositions of *Trachurus trachurus*: A fish commonly consumed in Nigeria. *Journal of Emerging Trends in Engineering and Applied Sciences*, **4**(3): 378 – 385.
- Ahmed, R. S. and Sharma, S. B. (1997). Biochemical studies on combined effects of garlic (*Allium sativum* L) and ginger (*Zingiber officinale* R.) in albino rats. *Indian Journal of Experimental Biology*, **35**(8): 841 – 843.
- Ajay, M., Gilanui, A. H. and Mustafa, M. R. (2003). Effect of flavonoids on vascular smooth muscles of the isolated rat thoracic aorta. *Life Sciences*, **74**(5): 603-612.
- Akoachere, J. F., Ndip, R. N. and Chenwi, E. B. (2002). Antibacterial effect of *Zingiber officinale* and *Garcinia kola* on respiratory tract pathogens. *East African Medical Journal*, **79**(11): 88-92.
- Al-Awwadi, N. A. J. (2017). Potential health benefits and scientific review of ginger. *Journal of Pharmacognosy and Phytotherapy*, **9**(7): 111-116.
- Amagase, H., Petesch, B. L., Matsuura, H., Kasuga, S. and Itakura, Y. (2001). Intake of Garlic and Its Bioactive Components. *The Journal of Nutrition*, **131**(3): 955 – 962.
- Amagase, H. (2006). Clarifying the real bioactive constituents of garlic. *Journal of Nutrition*, **136**(3): 716 – 725.
- Dashak, D. A., Dawang, M. L. and Lucas, N. B. (2001). An assessment of the proximate composition of locally produced spices known as dadawa basso and dadawa kawla from three markets in Plateau State of Nigeria. *Food Chemistry*, **75**(2): 231-235.
- El-Baroty, G. S., Abd El-Baky, H. H., Farag, R. S. and Saleh M. A. (2010). Characterization of antioxidant and antimicrobial compounds of cinnamon and ginger essential oils. *African Journal of Biochemistry Research*, **4**(6): 167 – 174.
- Chinedu, I. and Jivini, S. Z. (2019). Medicinal Properties of Ginger and Garlic: A Review. *Current Trends in Biomedical Engineering and Biosciences*, **18**(2): 47 – 52.
- Chukwu, O. and Shaba, M. (2009). Effects of drying methods on proximate compositions of catfish (*Clarias gariepinus*). *World Journal of Agricultural Sciences*, **5**(1): 114 – 116.
- El-Lahamy, A. A., Khalil, K. I., El-Sherif, S. A., Ibrahim, H. R. and Mahmud, A. A. (2019). Changes in fish during cooking methods (frying and grilling): A review. *Journal of Public Health Catalog* **2**(2): 169 – 172.
- Eyo, A. (2001). Shelf-life of moon fish (*Citharinus citharinus*): Drying storage at ambient temperature F.A.O fisheries report No 574: 35-37.
- Fenwick, G. R. and Hanley, A. B. (1985). The genus allium. Part 2. *Critical Review in Food Science and Nutrition*, **22**: 273 – 377.
- Gana, A. B., Ibrahim, R., Iriobe, T., Jega, I. S. and Mohammed, K. S. (2020). Proximate and mineral compositions of different fish species from Yamama Lake in Kalgo Local Government Area of Kebbi State, Nigeria. *Journal of Agriculture and Food Environment*, **7**(2): 17 – 25.
- Gebreselema, G. and Mebrahtu, G. (2013). Medicinal values of garlic: A review. *International Journal of Medicine and Medical Sciences*, **5**(9): 401 – 408.
- Hassan, L. G., and Umar, K. J. (2004). Proximate and mineral composition of seeds and pulp of African locust bean (*Parkia biglobosa* L.). *Nigerian Journal of basic and applied sciences*, **13**: 15-27.
- Hosseini, H., Mahmoudzadeh, M., Rezaei, M., Mahmoudzadeh, L., Khaksar, R., Karimian-Khosroshahi, N. and Babakhani, A. (2014). Effect of different cooking methods on minerals, vitamins and nutritional quality indices of kutum roach (*Rutilus frisii kutum*). *Food Chemistry*, **148**: 86 – 91.

- Ibrahim, R., Gana, A. B., Iriobe, T. and Hassan, S. A. (2018). Effect of Different Processing Methods on the Chemical Composition of *Mangifera Indica* Leaves. *Journal of Agriculture and Environment*, **14**(2): 201 – 207.
- Idris, G.L., Omojowo, F.S., Omojasola, P.F., Adetunji, C.O. and Ngwu, E.O. (2010). The effect of different concentrations of ginger on the quality of smoked dried catfish (*Clarias gariepinus*). *Nature and Science*, **8**(4): 59 – 63.
- Iheagwara, M. C. (2013). Effect of ginger extract on stability and sensorial quality of smoked mackerel (*Scomber scombrus*) fish. *Journal of Nutrition and Food Sciences*, **3**(3): 1 – 5.
- Karimian-Khosroshahi, N., Hosseini, H., Rezaei, M., Khaksar, R. and Mahmoudzadeh, M. (2016). Effect of different cooking methods on minerals, vitamins, and nutritional quality indices of rainbow trout (*Oncorhynchus mykiss*). *International Journal of Food Properties*, **9**(11): 2471 – 2480.
- Lanzotti, V., Scala, F., and Bonanomi, G. (2014). Compounds from *Allium* species with cytotoxic and antimicrobial activity. *Phytochemistry Reviews*, **13**: 769 – 791.
- Lawson, L. D. (1993). Bioactive organosulfur compounds of garlic and garlic products. In: ACS Symposium Series 534, Human Medicinal Agents from Plants (Kinghorn, A. D. & Balandrin, M. F., eds.). *American Chemical Society*, 306 – 330.
- Mader, F. H. (1990). Treatment of hyperlipidaemia with garlic-powder tablets. Evidence from the German Association of General Practitioners' multicentric placebo-controlled double-blind study. *Arzneimittelforschung*, **40**(10): 1111 – 1116.
- Musa, U., Hati, S. S., Adamu, Y. I. and Mustapha, A. (2010). Pesticides Residues in Smoked Fish Samples from North-Eastern Nigeria. *Journal of Applied Sciences*, **10**: 975 – 980.
- Nwinuka, N. M., Ibeh, G. O. and Ekeke, G. I. (2005). Proximate Composition and Levels of Some Toxicants in Four Commonly Consumed Spices. *Journal of Applied Sciences and Environmental Management*, **9**(1): 150 – 155.
- Ohaeri, O. C. and Adoga G. I. (1998). Effects of garlic oil on body fluid acidity and electrolytes in experimental diabetic rats. *Medical Science Research*, **26**: 487.
- Olaosebikan, B. D. and Raji, A. (2004). *A Field Guide to Nigerian Freshwater Fishes*. 2nd Edition. Federal College of Freshwater Fisheries Technology, New Bussa, Niger State, Nigeria.
- Olopade, O. A., Taiwo, I. O. and Agbato, D. A. (2013). Effect of traditional smoking method on nutritive values and organoleptic properties of *Sarotherodon galilaeus* and *Oreochromis niloticus*. *International Journal of Applied Agricultural and Apicultural Research*, **9**(1&2): 91 – 97.
- Omitoyin, B. O. (2007). *Introduction to Fish Farming in Nigeria*. Ibadan University Press, University of Ibadan. ISBN: 9781214279. 105pp.
- Onyia, L. U and Adebayo, E. F. (2008). Fish Processing and Marketing Livelihood Opportunity for women in Adamawa State *Nigerian Journal of Tropical Agriculture*, **10**: 33 – 39.
- Otunola, G. A., Oloyede, O. B., Oladiji, A. T. and Afolayan, A. J. (2010). Comparative analysis of the chemical composition of three spices – *Allium sativum* L. *Zingiber officinale* Rosc. and *Capsicum frutescens* L. commonly consumed in Nigeria. *African Journal of Biotechnology*, **9**(41): 6927 – 6931.
- Papu, S., Jaivir, S., Sweta, S. and Singh, B. R. (2017). Medicinal values of Garlic (*Allium sativum* L.) in Human Life: An Overview. *Greener Journal of Agricultural Sciences*, **4**(6): 265 – 280.
- Park, Y. J., Wen, J., Bang, S., Park, S. W. and Song, S. Y. (2006). 6-gingerol induces cell cycle arrest and cell death of mutant p53-expressing pancreatic cancer cells. *Yonsei Medical Journal*, **47**(5): 688 – 697.
- Pingxiang, L., Rui, W., Yanyang X., Yue, F., Linjuan H., Yongzhong, Q. and Jing, Q. (2020). Metabolic Changes in Different Tissues of Garlic Plant during Growth. *Journal of Agricultural and Food Chemistry*, **68**(44): 12467 – 12475.

- Ranilla, L. G., Kwon, Y. I., Apostolidis, C. and Shetty, K. (2010). Phenolic compounds, antioxidant activity and *in-vitro* inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly used medicinal plants, herbs and spices in Latin America. *Bioresource Technology*, **101**(12): 4676 – 4689.
- Rodrigo, A., Saray Q., Rocío, I. L., Enrique, O. F., Juan, P. R., Lucrecia, C. and Daniel, O. (2015). Immunomodulation and Anti-Inflammatory Effects of Garlic Compounds. *Journal of Immunology Research*, **3**: 1 – 13.
- Sampaio, R. M., Marcos, S. K., Moraes, I. C. F. and Perez, V. H. (2009). Moisture adsorption behaviour of biscuits formulated using wheat, oatmeal and passion fruit flour. *Journal of Food Processing and Preservation*, **33**(1): 105 – 113.
- Siegers, C. P. (1992). *Allium sativum*. In: Adverse Effects of Herbal Drugs (De Smet, P.A.G.M., Keller, K, Ha' Nsel, R. & Chandler, R. F., eds.), Springer-Verlag, Berlin, Germany, 73 – 77.
- Singh, D. (2013). Phenology of selected tree and shrubs species found in Aliero Local Government Area, Kebbi State, Nigeria. *Equity Journal of Sciences and Technology*, **1**(1): 23 – 26.
- Singh, R. N., Prithviraj B., Singh U. P., Singh, P. K. and Wagner, K. G. (1996). Effect of some pesticides and ajoene on biology of *Tetranychus urticae* Koch. *Journal of Plant Diseases and Protection*, **103**(2): 195 – 199.
- Srivastava, A., Shukla, Y. N. and Kumar, S. (2000). Recent development in plant derived antimicrobial constituents A Review. *Journal of Medical and Aromatic Plant Sciences*, **22**(2): 349 – 405.
- Stephen, N. M., Shakila, R. J., Jeyasekaran, G. and Sukumar, D. (2010). Effect of different types of heat processing on chemical changes in tuna. *Journal of Food Science Technology*, **47**(2): 174 – 181.
- Suetsuna, K. (1998). Isolation and characterization of angiotensin I-converting enzyme inhibitor dipeptides derived from *Allium sativum* (garlic). *The Journal of Nutritional Biochemistry*, **9**(7): 415 – 419.
- Sumiyoshi, H. (1997). New pharmacological activities of garlic and its constituents. *Nippon Yakurigaku Zasshi*, **110**(1): 93 – 97.
- Synnes, M., Larseen W. E., and Kjerstad, M. (2007). Chemical characterization and properties of five deep-sea fish species. *LWT-Food Sci Technol* **40**: 1049-1055.
- Wang, B. H., Zuzel, K. A., Rahman, K. and Billington, D. (1998). Protective effects of aged garlic extract against bromobenzene toxicity to precision-cut rat liver slices. *Toxicology*, **126**(3): 213 – 222.
- Yakubu, M. M., and Ngueku, B. B. (2015). Quality assessment of smoked-dried fish from five different markets in Lafia, Nigeria. *International Journal of Fisheries and Aquatic Studies*, **2**(4): 135-139.
- Yin, M. C. and Cheng, W. S. (2003). Antioxidant and antimicrobial effects of four garlic-derived organosulfur compounds in ground beef. *Meat Sciences*, **63**: 23 – 28.
- Yoshida, H., Iwata, N., Katsuzaki, H., Naganawa, R., Ishikawa, K., Fakuda, H., Fujino T. and Suzuki, A. (1998). Antimicrobial activity of a compound isolated from an oil-macerated garlic extract. *Bioscience, Biotechnology and Biochemistry*, **62**(5): 1014 – 1017.
- Young, H. Y., Luo, Y. L., Cheng, H. Y., Hsieh, W. C., Liao, J. C. and Peng, W. H. (2005). Analgesic and anti-inflammatory activities of (6)-gingerol. *Journal of Ethnopharmacology*, **96**(1): 207 – 210.
- Zafar, F. H. S., Zahid, M. and Bat, L. (2019). The effects of traditional frying method on proximate composition and energetic values of fish species from Karachi coast of Pakistan. *Korean Journal of Food and Health Convergence*, **5**(2): 35 – 43.

Table 1: Mean proximate and mineral compositions of ginger and garlic

Proximate composition (%)	Ginger	Garlic
Crude protein	5.69	9.19
Moisture content	4.50	6.00
Ash content	3.50	2.50
Ether extract	7.50	6.50
Crude fibre	1.50	0.50
Nitrogen free extract	77.31	75.31
Mineral composition (mg/kg)		
Sodium	70.00	85.00
Potassium	1200.00	1750.00
Calcium	0.75	0.66
Magnesium	1.60	1.45
Phosphorus	3.64	3.07

Table 2: Proximate composition of differently processed *Oreochromis niloticus* at week zero (0)

Treatments	Parameters (%)					
	CP	MC	AC	EE	CF	NFE
T1	34.68±0.04 ^c	4.16±0.14 ^a	32.05±0.03 ^a	9.53±0.02 ^{ef}	4.35±0.15 ^{ab}	15.54±0.21 ^d
T2	38.78±0.06 ^a	3.10±0.06 ^c	14.09±0.07 ⁱ	11.52±0.03 ^a	3.55±0.04 ^c	28.96±0.15 ^{ab}
T3	35.34±0.24 ^{bc}	3.55±0.06 ^b	26.12±0.07 ^b	10.15±0.10 ^d	4.13±0.07 ^b	20.71±0.40 ^c
T4	34.88±0.04 ^{bc}	3.50±0.05 ^b	24.73±0.12 ^e	9.61±0.06 ^e	4.13±0.07 ^b	23.15±0.26 ^c
T5	34.82±3.30 ^{bc}	3.17±0.12 ^c	15.64±0.08 ^g	11.06±0.03 ^b	3.55±0.32 ^c	31.75±3.16 ^a
T6	34.63±0.04 ^c	3.07±0.03 ^c	25.63±0.09 ^c	10.51±0.04 ^c	3.05±0.02 ^d	23.12±0.16 ^c
T7	35.36±0.05 ^{bc}	3.52±0.03 ^b	25.09±0.06 ^d	9.27±0.25 ^{fg}	4.58±0.04 ^a	22.18±0.35 ^c
T8	39.61±0.04 ^a	3.60±0.06 ^b	14.69±0.11 ^h	11.57±0.04 ^a	3.34±0.29 ^{cd}	27.19±0.42 ^b
T9	37.92±0.02 ^{ab}	3.13±0.08 ^c	14.10±0.06 ⁱ	11.24±0.19 ^{ab}	3.57±0.04 ^c	30.04±0.40 ^{ab}
T10	34.96±0.02 ^{bc}	3.62±0.08 ^b	25.36±0.22 ^{cd}	9.17±0.12 ^g	4.45±0.07 ^{ab}	22.43±0.34 ^c
T11	37.55±0.10 ^{abc}	3.10±0.06 ^c	16.16±0.12 ^f	10.56±0.05 ^c	3.28±0.25 ^{cd}	29.35±0.53 ^{ab}
T12	35.14±0.03 ^{bc}	3.08±0.05 ^c	24.60±0.08 ^e	9.14±0.10 ^g	4.62±0.08 ^a	23.42±0.33 ^c
Proximate compositions after eight (8) weeks of storage						
T1	34.34±0.03 ^a	4.54±0.02 ^c	29.04±0.02 ^d	11.04±0.02 ^f	2.55±0.03 ^d	18.49±0.11 ^g
T2	26.09±0.01 ^h	2.55±0.03 ^g	21.06±0.03 ^k	28.56±0.03 ^a	5.55±0.03 ^a	16.19±0.11 ^h
T3	23.56±0.01 ^j	4.56±0.03 ^c	31.04±0.02 ^a	9.04±0.03 ^h	1.55±0.03 ^e	30.25±0.11 ^a
T4	28.37±0.03 ^e	3.05±0.03 ^f	26.54±0.02 ^g	11.06±0.03 ^f	3.04±0.02 ^c	27.93±0.11 ^b
T5	25.30±0.01 ⁱ	3.04±0.03 ^f	18.54±0.02 ^l	27.55±0.03 ^c	5.56±0.03 ^a	20.01±0.11 ^f
T6	29.45±0.03 ^c	6.02±0.02 ^a	29.54±0.02 ^c	11.04±0.02 ^f	2.53±0.02 ^d	21.41±0.11 ^e
T7	23.38±0.01 ^k	4.54±0.02 ^c	30.04±0.02 ^b	9.03±0.02 ^h	2.52±0.01 ^d	30.49±0.08 ^a
T8	26.09±0.01 ^h	4.53±0.03 ^c	23.54±0.02 ^h	27.55±0.02 ^c	5.53±0.02 ^a	12.76±0.10 ^j
T9	27.67±0.01 ^f	5.03±0.02 ^b	28.55±0.03 ^e	10.55±0.02 ^g	1.53±0.02 ^e	26.68±0.09 ^c
T10	28.75±0.03 ^d	4.04±0.03 ^d	27.03±0.02 ^f	11.55±0.02 ^e	2.54±0.02 ^d	26.08±0.11 ^d
T11	26.37±0.02 ^g	3.54±0.02 ^e	21.55±0.03 ^j	28.05±0.03 ^b	5.56±0.03 ^a	14.94±0.12 ⁱ
T12	31.07±0.01 ^b	4.55±0.03 ^c	23.04±0.02 ⁱ	27.05±0.03 ^d	4.04±0.03 ^b	10.25±0.11 ^k

Mean with the same superscript along the same column are not significantly different ($P \geq 0.05$), T1 (oven drying), T2 (Frying), T3 (smoking), T4 (oven dried + 5% ginger-garlic), T5 (Fried + 5% ginger-garlic), T6 (smoked + 5% ginger-garlic), T7 (oven dried + 10% ginger-garlic), T8 (Fried + 10% ginger-garlic), T9 (smoked + 10% ginger-garlic), T10 (oven dried + 15% ginger-garlic), T11 (Fried + 15% ginger-garlic), T12 (smoked + 15% ginger-garlic), CP (crude protein) MC (moisture content) AC (ash content), EE (ether extract), CF (crude fibre) NFE (nitrogen free extract)

Table 3: Mineral composition of differently processed *Oreochromis niloticus* at week zero (0)

Treatments	Parameters (mg/kg)				
	Sodium	Potassium	Calcium	Magnesium	Phosphorus
T1	655.00±2.89 ^c	1405.00±2.89 ^c	1.18±0.01 ^d	2.56±0.03 ^a	4.04±0.03 ^{ab}
T2	405.00±2.89 ^f	1304.67±2.60 ^d	1.27±0.01 ^b	2.44±0.02 ^b	3.57±0.02 ^e
T3	705.00±2.89 ^b	1450.00±2.89 ^b	1.14±0.02 ^d	2.57±0.01 ^a	3.99±0.01 ^c
T4	700.00±2.89 ^b	1500.00±2.89 ^a	1.07±0.01 ^e	2.48±0.01 ^b	4.05±0.01 ^{ab}
T5	451.67±1.67 ^e	1250.00±2.89 ^e	1.35±0.03 ^a	2.34±0.02 ^{cd}	3.40±0.01 ^e
T6	655.00±2.89 ^c	1405.00±2.89 ^c	1.15±0.01 ^d	2.57±0.01 ^a	3.98±0.01 ^c
T7	755.00±2.89 ^a	1450.00±2.89 ^b	1.07±0.01 ^e	2.47±0.01 ^b	4.03±0.01 ^{ab}
T8	350.00±2.89 ^g	1550.00±2.89 ^e	1.25±0.03 ^{bc}	2.27±0.01 ^d	3.57±0.02 ^e
T9	411.67±6.01 ^f	1216.00±8.82 ^f	1.29±0.02 ^{ab}	2.18±0.02 ^e	3.69±0.01 ^d
T10	588.33±30.87 ^d	1210.00±5.77 ^f	1.07±0.01 ^e	2.57±0.04 ^a	4.07±0.02 ^a
T11	476.67±14.53 ^e	1408.00±4.41 ^c	1.15±0.03 ^d	2.36±0.03 ^c	3.59±0.02 ^e
T12	758.33±4.41 ^a	1455.00±2.89 ^b	1.19±0.02 ^{cd}	2.48±0.01 ^b	4.02±0.04 ^{ab}
Mineral compositions after eight (8) weeks of storage					
T1	233.33±3.33 ^a	750.00±2.89 ^c	1.66±0.03 ^a	1.17±0.01 ^e	6.52±0.03 ^{ab}
T2	182.37±0.19 ^e	654.00±2.08 ^e	0.84±0.03 ^c	1.33±0.02 ^{bc}	4.17±0.02 ^g
T3	187.55±0.03 ^d	752.00±1.53 ^c	0.77±0.01 ^d	1.37±0.01 ^b	6.48±0.01 ^b
T4	192.53±0.01 ^c	901.67±1.67 ^a	0.85±0.02 ^c	1.45±0.03 ^a	5.29±0.02 ^d
T5	182.67±1.45 ^e	700.00±2.89 ^d	0.77±0.01 ^d	1.28±0.02 ^{cd}	5.27±0.02 ^d
T6	207.20±0.15 ^b	805.00±2.89 ^b	1.48±0.01 ^b	1.17±0.01 ^e	4.48±0.01 ^e
T7	172.58±0.06 ^f	550.00±28.87 ^f	0.68±0.01 ^e	1.16±0.02 ^e	4.06±0.02 ^h
T8	182.56±0.03 ^e	701.67±1.67 ^d	0.85±0.03 ^c	1.37±0.01 ^b	4.24±0.02 ^f
T9	181.67±1.67 ^e	704.33±2.33 ^d	0.73±0.01 ^{de}	1.25±0.03 ^d	5.26±0.02 ^d
T10	182.50±0.29 ^e	651.67±1.67 ^e	0.77±0.01 ^d	1.37±0.01 ^b	6.57±0.01 ^a
T11	187.55±0.03 ^d	751.67±1.67 ^c	0.87±0.01 ^c	1.44±0.02 ^a	6.48±0.01 ^b
T12	180.18±0.18 ^e	650.00±2.89 ^e	0.77±0.01 ^d	1.43±0.01 ^a	5.59±0.01 ^c

Mean with different superscript along the same column are significantly different ($P \leq 0.05$), T1 (oven drying), T2 (Frying), T3 (smoking), T4 (oven dried + 5% ginger-garlic), T5 (Fried + 5% ginger-garlic), T6 (smoked + 5% ginger-garlic), T7 (oven dried + 10% ginger-garlic), T8 (Fried + 10% ginger-garlic), T9 (smoked + 10% ginger-garlic), T10 (oven dried + 15% ginger-garlic), T11 (Fried + 15% ginger-garlic), T12 (smoked + 15% ginger-garlic)